

Longitudinal Oscillations in a Nonextensive Relativistic Plasma

Víctor Muñoz

*Department of Earth System Science and Technology,
Kyushu University, Fukuoka 816-8580, Japan and*

Departamento de Física, Facultad de Ciencias, Universidad de Chile, Casilla 653, Santiago, Chile

The dispersion relation of longitudinal electrostatic oscillations in a relativistic plasma is studied in the context of the nonextensive statistics formalism proposed by Tsallis [C. Tsallis, J. Stat. Phys. **52**, 479 (1988)], where nonextensivity is characterized by a parameter q in Tsallis's entropy. $q = 1$ corresponds to the usual Boltzmann-Gibbs, extensive statistics formalism. In the nonrelativistic regime, normalizability of the equilibrium distribution function implies that $-1 \leq q \leq \infty$. We show that in the relativistic regime much tighter constraints must be satisfied, namely $0 \leq q \leq 1 + k_B T / mc^2$, where k_B is the Boltzmann constant, T is the temperature of the plasma, and m is the particle mass.

Then we study longitudinal oscillations in a proton-electron plasma, assuming immobile protons, and electrons whose distribution function maximizes Tsallis's entropy. The dispersion relation of these oscillations is written in integral form for the long wavelength limit. Explicit expressions in terms of generalized hypergeometric functions can be found for all possible values of q in the ultra-relativistic regime.

I. INTRODUCTION

Traditionally, the equilibrium of statistical systems has been studied based on the Boltzmann-Gibbs entropy,

$$S_{BG} = -k_B \sum_i p_i \ln p_i, \quad (1)$$

where k_B is the Boltzmann constant, and p_i is the probability of the i -th microstate. However, an increasing amount of experimental, computational and theoretical evidence, shows that this formalism is not adequate for systems where long range interactions in time or space are important, and, in general, systems where the relevant phase space has fractal or multifractal structure. Tsallis [1] proposed the following form for the entropy of such systems:

$$S_q = k_B \frac{1 - \sum_i p_i^q}{q - 1}, \quad (2)$$

where q is a real number. For $q = 1$, S_q reduces to S_{BG} .

S_q has the basic property that for independent systems A and B , the entropy of the composite system $A + B$ is given by:

$$S_q(A + B) = S_q(A) + S_q(B) + (1 - q)S_q(A)S_q(B). \quad (3)$$

Thus, q is a measure of the nonextensivity. Among the various physical systems where connections with the Tsallis entropy formalism have been found are gravitational systems [2], long range Hamiltonian systems [3], nonlinear inverse bremsstrahlung absorption in plasmas [4], turbulence [5], and granular systems [6], and many applications in astrophysics, fluid dynamics, biology, economy, quantum mechanics, and others. (See, e.g. [7].)

An important property of the nonextensive formalism is that the distribution function which maximizes S_q is non-Maxwellian [8, 9]. Specifically, for $q < 1$, high energy states are more probable than in the extensive case; for $q > 1$ high energy states are less probable than in the extensive case, and there is a cutoff beyond which no states exist. For a unidimensional gas where $E = mv^2/2$, this cutoff is given by [9]

$$v_{\max} = \sqrt{2k_B T / m(q - 1)}. \quad (4)$$

Velocity distributions in plasmas are often found to be non-Maxwellian. High energy tails in space and laboratory plasmas [10, 11, 12, 13, 14] are a common occurrence. Metaequilibrium non-Maxwellian radial density profiles in cylindrical turbulent pure electron plasmas are observed [15]. As stated above, such distribution functions emerge naturally if the system is described by a nonextensive statistics, and several authors have successfully applied this approach to plasma systems [4, 16, 17, 18].

Lima et al. [9] studied the longitudinal oscillations of a collisionless thermal plasma, in the context of the nonextensive formalism based on (2). They derived the dispersion relation for electrostatic waves, and calculated it explicitly

for an equilibrium distribution function which maximizes (2). They found that it fits experimental data better than if the energy distribution is given by a Maxwellian.

All the previous analysis have been made for non-relativistic plasmas. However, relativistic extensions of the Tsallis formalism are possible [19]. Our purpose is to study the consequences of a relativistic nonextensive statistics in plasmas. In particular, we will examine some general properties of the equilibrium distribution function and calculate the dispersion relation for electrostatic waves in the long wavelength limit for ultrarelativistic plasmas.

II. RELATIVISTIC DISTRIBUTION FUNCTION

In the nonextensive description, the equilibrium distribution function for a relativistic plasma can be written [19]

$$f_q(\vec{p}) = A_q \left[1 - (q-1) \frac{E}{k_B T} \right]^{1/(q-1)}, \quad (5)$$

where

$$E = \sqrt{m^2 c^4 + \vec{p}^2 c^2}, \quad (6)$$

is the particle energy and A_q is a normalization constant.

Equation (5) generalizes the nonrelativistic expression (see, e.g., [9]). For $q = 1$, (5) yields the usual Boltzmann distribution function $f_{q=1} = A_1 \exp(-E/k_B T)$.

For simplicity, in the following we will consider a one dimensional plasma, so that $\vec{p} = p\hat{z}$.

Since the relativistic energy is positive, f_q will be real and normalizable if E satisfies the following conditions:

$$\text{if } q \geq 1 \quad 0 \leq E \leq \frac{k_B T}{q-1}, \quad (7)$$

$$\text{if } q \leq 1 \quad 0 \leq E, \quad (8)$$

i.e., E can take any possible value if $q \leq 1$, but the distribution function has an energy cutoff if $q \geq 1$. Notice that both (7) and (8) hold for the extensive limit $q = 1$, in which case they give the expected constraint $0 \leq E < \infty$.

Due to the cutoff (7), if $q \geq 1$ f_q is always normalizable. However, normalizability of the distribution function for $0 \leq E \leq \infty$ when $q < 1$ is not guaranteed. The integral

$$I = \int_{-\infty}^{\infty} f_q(p) dp, \quad (9)$$

is finite if $f_q(p) \sim p^s$ when $p \rightarrow \infty$, with $s < -1$. Using (6),

$$f_q(p) \xrightarrow{p \rightarrow \infty} C p^{\frac{1}{q-1}},$$

where $C = \left[-\frac{q-1}{k_B T} \right]^{\frac{1}{q-1}} > 0$. Thus, $f_q(p)$ is normalizable if $q > 0$.

Let us now analyze the case $q \geq 1$. The cutoff in (7) and (6) give

$$|p| \leq \frac{1}{c} \sqrt{\left(\frac{k_B T}{(q-1)} \right)^2 - m^2 c^4}. \quad (10)$$

If

$$k_B T / (q-1) \geq m c^2, \quad (11)$$

then (10) yields the cutoff in momentum space corresponding to the energy cutoff (7). In the limit case $k_B T / (q-1) = m c^2$, the only momentum allowed is $p = 0$. If $k_B T / (q-1) < m c^2$, then no real value of p satisfies condition (7), and no distribution function exists. We may interpret this saying that for a given temperature, only q values consistent with (11) are possible:

$$q \leq 1 + \frac{k_B T}{m c^2}. \quad (12)$$

Summarizing:

Case 1: $0 \leq q$.

Forbidden q . f_q is not normalizable.

Case 2: $0 < q \leq 1$.

f_q is normalizable. Energy can take values

$$0 \leq E < \infty .$$

Case 3: $1 \leq q \leq 1 + \frac{k_B T}{mc^2}$.

f_q is normalizable. Energy can only take values

$$0 \leq E \leq \frac{k_B T}{q-1} ,$$

i.e.

$$0 \leq |p| \leq \frac{1}{c} \sqrt{\left(\frac{k_B T}{(q-1)} \right)^2 - m^2 c^4} .$$

Case 4: $1 + \frac{k_B T}{mc^2} < q$.

Forbidden q . No values of p consistent with energy cutoff.

Two main differences arise with respect to the non-relativistic case [9], where allowed values are $-1 \leq q < \infty$. In the relativistic case constraints are much tighter, $0 \leq q \leq 1 + k_B T/mc^2$, and depend on the ratio of thermal to rest energy.

Considering these constraints, the relativistic distribution function can be plotted. The result is qualitatively similar to the nonrelativistic case:




FIG. 1: Relativistic dispersion function (5) ($k_B T = m = c = 1$). Full circles: $q = 0.5$. Solid line: $q = 1$. Open circles: $q = 1.5$.

III. DISPERSION RELATION FOR LONGITUDINAL PLASMA OSCILLATIONS

We consider a proton-electron relativistic plasma. Assuming protons remain at rest, and that particles only move in one dimension ($\hat{z}$), the dispersion relation for longitudinal oscillations is

$$1 = i \frac{\omega_p^2}{k} m \int dp_z \frac{\partial f_q(p_z)/\partial p_z}{s + i k v_z} , \quad (13)$$

where ω_p is the plasma frequency, k is the wavenumber, s is the argument of the Laplace transform ($s = -i\omega$, where ω is the wave frequency, if there is no damping), m the electron mass, $p = m\gamma v$ is the particle momentum, and $f_q(p, z)$ is the equilibrium electron distribution function, normalized to unity. Particle energy is $E = \sqrt{m^2 c^4 + p_z^2 c^2}$.

Following Ref. [9], we now consider the long wavelength limit $k \ll k_D$, where $k_D = 2\pi/\lambda_D$, and λ_D is the Debye length. The integrand has a pole at $v = -is/k$. If $\text{Im } v \ll 1$, integration can be performed on the real axis:

$$1 \simeq \frac{im\omega_p^2}{sk} \int_{-p_{\max}}^{p_{\max}} dp_z \frac{\partial f_q}{\partial p_z} \left(1 - i \frac{kv_z}{s} + \dots \right), \quad (14)$$

where $p_{\max}$ is the maximum momentum allowed by (10) if $1 \leq q \leq 1 + k_B T/mc^2$, or ∞ if $0 < q \leq 1$. Since f_q is an even function of p_z , the first term in (14) vanishes. The dispersion relation can then be written:

$$1 = \frac{m\omega_p^2}{s^2} \int_{-p_{\max}}^{p_{\max}} dp_z \frac{p_z}{m\gamma} \frac{\partial f_q}{\partial p_z},$$

where $\gamma = (1 + p_z^2/m^2 c^2)^{1/2}$. Integrating by parts, and using the fact that $f_q(\pm p_{\max}) = 0$, finally yields

$$1 = -\frac{\omega_p^2}{s^2} \int_{-p_{\max}}^{p_{\max}} dp_z \frac{f_q(p_z)}{\gamma^3}. \quad (15)$$

Equation (15) depends on q in general, and therefore, unlike the nonrelativistic case [9], nonextensive corrections to the dispersion relation of longitudinal waves may appear to the lowest order in kv_z/s .

IV. ULTRARELATIVISTIC PLASMA

Closed expressions in terms of known functions can be obtained for Eq. (15) in the ultrarelativistic case $|p_z| \gg mc$. As shown in Sec. II, q can only take values between 0 and $1 + k_B T/mc^2$. We consider two situations:

A. $0 < q \leq 1$

Since $p_{\max} = \infty$, in this case, the dispersion relation (15) can be written

$$1 = -2 \frac{\omega_p^2}{s^2} A_q \int_0^\infty dp_z \frac{1}{\left(1 + \frac{p_z^2}{(mc)^2}\right)^{3/2}} \frac{1}{\left(1 - (q-1) \frac{p_z c}{k_B T}\right)^{-\frac{1}{q-1}}}. \quad (16)$$

Noting that $-(q-1) > 0$ and $-1/(q-1) > 0$, normalization of the distribution function

$$f_q(p_z) = A_q \frac{1}{\left(1 - (q-1) \frac{|p_z| c}{k_B T}\right)^{-\frac{1}{q-1}}}, \quad (17)$$

$$1 = \int_0^\infty \hat{f}_q(p_z) dp_z, \quad (18)$$

yields

$$A_q = \frac{1}{mc} \frac{q}{2\tau}, \quad (19)$$

where

$$\tau = \frac{k_B T}{mc^2}. \quad (20)$$

The integral in (16) can be done analytically, yielding

$$1 = -\frac{\omega_p^2}{s^2} \frac{1}{\tau} q I \left(\tau, -\frac{1}{q-1} \right), \quad (21)$$

where

$$\begin{aligned}
I(\tau, \alpha) = & -\frac{1}{\sqrt{\pi}}(-\tau\alpha)^{\alpha+1}\alpha\Gamma\left(\frac{3+\alpha}{2}\right)\Gamma\left(-\frac{\alpha}{2}\right){}_2F_1\left(\frac{1+\alpha}{2}, \frac{3+\alpha}{2}, \frac{3}{2}; -\alpha^2\tau^2\right) \\
& + \frac{1}{\sqrt{\pi}}(-\tau\alpha)^\alpha\Gamma\left(\frac{1-\alpha}{2}\right)\Gamma\left(\frac{2+\alpha}{2}\right){}_2F_1\left(\frac{2+\alpha}{2}, \frac{\alpha}{2}, \frac{1}{2}; -\alpha^2\tau^2\right) \\
& + \frac{\tau\alpha}{1-\alpha}{}_3F_2\left[\left\{\frac{1}{2}, 1, \frac{3}{2}\right\}, \left\{\frac{2-\alpha}{2}, \frac{3-\alpha}{2}\right\}, -\tau^2\alpha^2\right], \tag{22}
\end{aligned}$$

${}_2F_1$ is the hypergeometric function,

$${}_2F_1(a, b, c; z) = \sum_{\nu=0}^{\infty} \frac{(a)_\nu (b)_\nu}{(c)_\nu} \frac{z^\nu}{\nu!}, \tag{23}$$

with $(a)_\nu = a(a+1)\cdots(a+\nu-1)$, and ${}_mF_n$ is the generalized hypergeometric function,

$${}_mF_n(\{a_1, \dots, a_m\}, \{b_1, \dots, b_n\}, z) = \sum_{\nu=0}^{\infty} \frac{(a_1)_\nu \cdots (a_m)_\nu}{(b_1)_\nu \cdots (b_n)_\nu} \frac{z^\nu}{\nu!}. \tag{24}$$

$$\mathbf{B.} \quad 1 \leq q \leq 1 + k_B T / mc^2$$

Now $p_{\max}$ is given by Eq. (10), and dispersion relation Eq. (15) is

$$1 = -2 \frac{\omega_p^2}{s^2} A_q \int_0^{p_{\max}} dp_z \frac{1}{\left(1 + \frac{p_z^2}{(mc)^2}\right)^{3/2}} \left(1 - (q-1) \frac{p_z c}{k_B T}\right)^{\frac{1}{q-1}}. \tag{25}$$

Noting that $q-1 > 0$, normalization of the distribution function yields

$$A_q = \frac{q}{2mc\tau} \frac{1}{\tau} \left[1 - \left(1 - (q-1) \frac{\hat{p}}{\tau}\right)^{\frac{q}{q-1}}\right]^{-1}, \tag{26}$$

where

$$\hat{p} = \frac{p_{\max}}{mc}. \tag{27}$$

Again, integration in (25) can be performed analytically, yielding

$$1 = -\frac{\omega_p^2}{s^2} \frac{1}{\sqrt{1 + \hat{p}^2}} J\left(\tau, \frac{1}{q-1}\right), \tag{28}$$

where

$$\begin{aligned}
J(\tau, \alpha) = & \left[1 - \left(1 - \frac{\hat{p}}{\alpha\tau}\right)^{\alpha+1}\right]^{-1} \frac{1}{(\alpha^2\tau^2 + 1)^2} \\
& \times \left[(1 + i\alpha\tau)\sqrt{1 + \hat{p}^2} \cdot F_1\left(1 + \alpha; \frac{3}{2}, \frac{3}{2}; 2 + \alpha; \frac{\alpha\tau}{\alpha\tau + i}, \frac{\alpha\tau}{\alpha\tau - i}\right) \right. \\
& \left. - \left(1 - \frac{\hat{p}}{\alpha\tau}\right)^{1+\alpha} [(1 + i\alpha\tau) + \hat{p}(i + \alpha\tau)] \cdot F_1\left(1 + \alpha; \frac{3}{2}, \frac{3}{2}; 2 + \alpha; \frac{\alpha\tau - \hat{p}}{\alpha\tau + i}, \frac{\alpha\tau - \hat{p}}{\alpha\tau - i}\right)\right], \tag{29}
\end{aligned}$$

where F_1 is the Appell hypergeometric function of two variables:

$$F_1(a; b_1, b_2; c; x, y) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(a)_{m+n} (b_1)_m (b_2)_n}{m! n! (c)_{m+n}} x^m y^n. \tag{30}$$

V. CONCLUSIONS

Some consequences of describing a plasma based on a nonextensive statistical treatment have been discussed. In particular, the one dimensional plasma distribution function which maximizes Tsallis's entropy S_q has been considered. In the nonrelativistic case, normalizability of the distribution function constrains q to values greater than or equal to -1 . However, in the relativistic case a much shorter range is possible, and the maximum value of q depends on the ratio of temperature to particle rest energy.

Then we derive the dispersion relation for longitudinal oscillations in a plasma composed of electrons and rest protons. In the long wavelength limit, and unlike the nonrelativistic case, nonextensive corrections appear to the lowest order. Finally, in the ultra-relativistic regime, this dispersion relation is written explicitly in terms of generalized hypergeometric functions.

-
- [1] C. Tsallis, J. Stat. Phys. **52**, 479 (1988).
 - [2] A. Taruya and M. aki Sakagami, Phys. Rev. Lett. **90**, 181101 (2003).
 - [3] V. Latora, A. Rapisarda, and C. Tsallis, Phys. Rev. E **64**, 056134 (2001).
 - [4] C. Tsallis and A. M. C. de Souza, Phys. Lett. A **235**, 444 (1997).
 - [5] T. Arimitsu and N. Arimitsu, Phys. Rev. E **61**, 3237 (2000).
 - [6] F. Sattin, J. Phys. A **36**, 1583 (2003).
 - [7] S. Abe and Y. Okamoto, *Nonextensive Statistical Mechanics and its Applications* (Springer-Verlag, Berlin Heidelberg, 2001).
 - [8] R. S. Jr., A. R. Plastino, and J. Lima, Phys. Lett. A **249**, 401 (1998).
 - [9] J. A. S. Lima, R. Silva, Jr., and J. Santos, Phys. Rev. E **61**, 3260 (2000).
 - [10] B. H. Mauk, S. M. Krimigis, E. P. Keath, A. F. Cheng, T. P. Armstrong, L. J. Lanzerotti, G. Gloeckler, and D. C. Hamilton, J. Geophys. Res. **92**, 15283 (1987).
 - [11] J. D. Scudder, J. E. C. Sittler, and H. S. Bridge, J. Geophys. Res. **86**, 8157 (1981).
 - [12] E. Marsch, K.-H. Mhlhuser, R. Schwenn, H. Rosenbauer, W. Pilipp, and F. M. Neubauer, J. Geophys. Res. **87**, 52 (1982).
 - [13] M. R. Collier, D. C. Hamilton, G. Gloeckler, P. Bochsler, and R. B. Sheldon, Geophys. Res. Lett. **23**, 1191 (1996).
 - [14] J. M. Liu, J. S. D. Groot, J. P. Matte, T. W. Johnston, and R. P. Drake, Phys. Rev. Lett. **72**, 2717 (1994).
 - [15] X.-P. Huang and C. F. Driscoll, Phys. Rev. Lett. **72**, 2187 (1994).
 - [16] B. M. Boghosian, Phys. Rev. E **53**, 4754 (1996).
 - [17] C. Anteneodo and C. Tsallis, J. Mol. Liq. **71**, 255 (1997).
 - [18] M. P. Leubner, Phys. Plasmas **11**, 1308 (2004).
 - [19] A. Lavagno, Phys. Lett. A **301**, 13 (2002).